

Semantic segmentation of Atomic Force Microscopy images of calcite topography

Chiara Recalcati^{*1}, Martina Siena¹, Gianlorenzo Bussetti², Monica Riva¹, Lamberto Duò² and Alberto Guadagnini¹

¹ Politecnico di Milano, Dipartimento di Ingegneria Civile e Ambientale, Milano, Italy

² Politecnico di Milano, Dipartimento di Fisica, Milano, Italy


*Contact: chiara.recalcati@polimi.it



POLITECNICO
MILANO 1863

PROBLEM STATEMENT

- Mineral dissolution/precipitation reactions are critical in many natural and industrial contexts.
- High resolution imaging techniques allowing a direct observation of the surface during the reaction revealed that they involve different dissolution mechanisms caused by an uneven local distribution of surface energy due to inhomogeneities and defects in the mineral lattice.



Typical features in the crystal lattice

- step
- impurities
- screw dislocation
- line defect
- point defect

- We rely on a multimodal Gaussian model to capture the spatial heterogeneity of *in-situ* and real time topography measurements of calcite samples subject to dissolution at far from equilibrium conditions acquired via Atomic Force Microscopy (AFM). Each Gaussian mode is associated with a characteristic dissolution pattern.
- We resort to an imaging segmentation technique to cluster topography data into different regions depending on the dissolution mechanism. Stochastic model parameters are then inferred on the data of the regions.

METHODS











IMAGE SEMANTIC SEGMENTATION & RESULTS



REMARKS

- Image semantic segmentation resulted to be very accurate in subdividing the mineral regions associated with different dissolution patterns.
- GMM parameters inferred on segmented regions provide an accurate representation of the sample frequency distribution.
- After training, the network is able to segment input images instantaneously.
- Future developments include the introduction of a third pixel class to subdivide nucleated monolayer etch pits from terraces.

REFERENCES

- Tahmasebi, P., Kamrava, S., Bai, T., Sahimi, M. Machine learning in geo- and environmental sciences: from small to large scale. *Advances in Water Resources* 142 (2020). doi.org/10.1016/j.advwatres.2020.103619